

Quenched Chiral Behavior of Hadrons with Overlap Fermions ^{*} [†]

Shao-Jing Dong ^a, Terrence Draper ^a, Ivan Horváth ^a, Frank Lee ^{b c}, and Jianbo Zhang ^d

^aDepartment of Physics and Astronomy, University of Kentucky, Lexington, KY 40506, USA

^bCenter for Nuclear Studies, Dept. of Physics, George Washington Univ., Washington, DC 20052, USA

^cJefferson Lab, 12000 Jefferson Avenue, Newport News, VA 23606, USA

^dCSSM and Dept. of Physics and Math. Physics, Univ. of Adelaide, Adelaide, SA 5005, Australia

We study the quenched chiral behavior of hadrons with the pseudoscalar mass as low as ≈ 280 MeV. We look for quenched chiral logs in the pion mass, determine the renormalized quark mass, and observe quenched artifacts in the a_0 and N^* propagators. The calculation is done on a quenched lattice of size 20^4 and $a = 0.148(2)$ fm using overlap fermions and an improved gauge action.

1. Simulation Details

Using a $\beta = 7.60$ (tree-level tadpole-improved) Lüscher–Weisz [1,2] gauge action, we study the chiral properties of hadrons on a 20^4 lattice with the overlap fermion [3,4]. The massive Dirac operator [5–7] is defined so that the tree-level renormalization of mass and wavefunction is unity.

$$\begin{aligned} D(m_0) &= (1 - \frac{m_0 a}{2\rho}) \rho D(\rho) + m_0 a \\ &= (\rho + \frac{m_0 a}{2}) + (\rho - \frac{m_0 a}{2}) \gamma_5 \epsilon(H) \end{aligned} \quad (1)$$

$\epsilon(H) = H/\sqrt{H^2}$, $H = \gamma_5 D_w$, and D_w is the usual Wilson fermion operator, except with a negative mass parameter $-\rho = 1/2\kappa - 4$ in which $\kappa_c < \kappa < 0.25$; we take $\kappa = 0.19$ in our calculation which corresponds to $\rho = 1.368$. See [8] for more details of the simulation.

2. Pion Mass and Chiral Logs

To search for chiral logs, we fit $m_\pi^2 a^2$ to [9]

$$m_\pi^2 a^2 = Am_0 a \{1 - \delta [\ln(Am_0 a / \Lambda_\chi^2 a^2) + 1]\} + Bm_0^2 a^2 \quad (2)$$

^{*}Presented by T. Draper at Lattice 2001, Berlin, Germany.

[†]This work is supported in part by the U.S. Department of Energy under grant numbers DE-FG05-84ER40154 and DE-FG02-95ER40907.

The best fits which give stable values of A and B and with errors less than half of the fitted values of δ for a range of $\Lambda_\chi = 0.6$ GeV to 1.4 GeV are listed in the table.

Table 1
Quenched chiral log parameter δ and χ^2/NDF as fitted from m_π^2 in Eq. (2).

Λ_χ	A	B	δ	χ^2/NDF
0.6	1.72(7)	3.0(8)	0.23(7)	0.18
0.8	1.42(11)	3.0(8)	0.28(11)	0.18
1.0	1.17(24)	3.0(8)	0.34(17)	0.18

We plot the fit with $\Lambda_\chi = 0.8$ GeV as a solid line in Fig. 1. To check if there is indeed a quenched chiral log, we fit $m_\pi^2 a^2$ alternatively without the chiral log term and find

$$m_\pi^2 a^2 = 1.88(10) m_0 a - 2.3(11) m_0^2 a^2 + 7.8(30) m_0^3 a^3 \quad (3)$$

with $\chi^2/NDF = 0.46$. A quadratic fit (shown in Fig. 1 as the dotted line) has $\chi^2/NDF = 0.90$. Thus the presence of a chiral log term is favored.

3. Quark Masses

To renormalize our estimates of the quark mass, we follow [7] to determine the renormalization constants $Z_M^{-1} = Z_S = Z_P$ and thus




Figure 1. $m_\pi^2 a^2$ versus $m_0 a$ calculated from $G_{A4P}(\vec{p}=0, t)$ with linear plus quadratic fit (dotted line) and chiral log fit (solid line) from Eq. (2) with $\Lambda_\chi = 0.8 \text{ GeV}$.

$m^{RGI} = Z_M(g_0)m_0(g_0)$: calculate

$$Z_M(g_0) = U_M \cdot \frac{1}{(r_0 m_0)} \Big|_{(r_0 m_\pi)^2 = x_{\text{ref}}} \quad (4)$$

using their tabulated values

$$U_M = \begin{cases} 0.181(6), & x_{\text{ref}} = 1.5736 \\ 0.349(9), & x_{\text{ref}} = 3.0 \\ 0.580(12), & x_{\text{ref}} = 5.0 \end{cases} \quad (5)$$

where U_M is a universal factor which they estimated from published results of another regularization, namely, $\mathcal{O}(a)$ -improved Wilson.

The value of the bare quark mass $m_0 a$ that reproduces a given pseudoscalar mass (through $x_{\text{ref}} = (r_0 m_\pi)^2$, with $r_0/a = 4.05(2)$ for our lattice) is obtained by interpolation of our data for the pion mass. From this we obtain $Z_M^{-1} = Z_S$ at three values of quark mass (the lowest two quark masses are roughly half-strange and strange) as shown in Fig. 2.

The operators and action are $\mathcal{O}(a)$ improved. For the axial vector renormalization constant, Z_A , we find small $(m_0 a)^2$ dependence as well [8,




Figure 2. Renormalization constant $Z_M^{-1} = Z_S$.

10]. Thus we fit $Z_M^{-1} = Z_S$ to a constant.

$$Z_M^{-1} = Z_S = 1.233(34) \quad \chi^2/NDF = 0.86 \quad (6)$$

From the $(m_\pi a)^2$ versus $m_0 a$ figure, we obtain $m_0 a$ at the physical pion mass. Using $m^{RGI} = m_0/Z_S$ and, from the 4-loop calculation with $N_f = 0$ [11], $Z_S^{MS} = Z_S/0.72076$, we obtain (very preliminary results) for $(m_u + m_d)/2$.

$$m^{RGI} = 7.3(7)(10) \text{ MeV} \quad (7)$$

$$m^{MS}(\mu = 2 \text{ GeV}) = 5.3(6)(7) \text{ MeV} \quad (8)$$

4. Chiral Loops in Quenched Propagators

Bardeen et al. [12] present evidence of a strong “quenched chiral loop” (QCL) effect, attributable to an $\eta' - \pi$ intermediate state, in the valence propagation of the scalar, isovector meson (a_0). The effect is manifested in a negative-metric contribution to the two-point correlation function which increases for decreasing quark mass. They were able to uncover the effect only after using their pole-shifting procedure, which mitigates the poor chiral properties of the Wilson fermions. We see this effect clearly with overlap fermions without the need for any such procedure.




Figure 3. 2pt corr. func. for a_0 with bare quark masses $m_0 a = 0.0150, 0.0164, 0.0192$, and 0.0274 .

Fig. 3 shows the two-point scalar-isovector local-local correlators for quark masses $m_0 a = 0.0150, 0.0164, 0.0192$, and 0.0274 ; they are plotted on a linear scale to show the correlator becoming negative at early times ($t/a \approx 1 - 2$, the source is at $t_0 = 3$) and then asymptotically approaching zero from below.

We also see the effect for $N^*(\frac{1}{2}^-)$. Fig. 4 shows the two-point N^* local-local correlator for quark masses $m_0 a = 0.0150, 0.0547$, and 0.1642 . The correlation function becomes negative at successively earlier times for decreasing mass. The effect becomes very pronounced at the lighter masses. See [13] for more details.

5. Summary

We found evidence for chiral logs in the dependence of the pion mass on quark mass. (See [8] for more details.) We also presented, for the first time, our very preliminary results for the renormalized light quark mass and the effect of chiral loops on the a_0 and N^* propagators. In a companion proceeding [10], we study zero modes and their effect on the pion propagator, and show our results for the axial renormalization constant Z_A and pion axial decay constant f_π , and the presence of chiral logs in the pseudoscalar decay constant, f_P . See [8] for complete details.




Figure 4. Two-point correlation function of N^* for bare quark masses $m_0 a = 0.0150, 0.0547$, and 0.1642 . Open (filled) symbols and bare masses labeled with a “(+)” (“(-)”) indicate positive (negative) values for the correlation function.

REFERENCES

1. M. Lüscher and P. Weisz, Phys. Lett. **B158**, 250 (1985).
2. M. Alford, W. Dimm, and P. Lepage, Phys. Lett. **B361**, 87 (1995).
3. H. Neuberger, Phys. Lett. **B417**, 141 (1998).
4. R. Narayanan and H. Neuberger, Nucl. Phys. **B443**, 305 (1995).
5. C. Alexandrou et al., Nucl. Phys. **B580**, 394 (2000), hep-lat/0002010.
6. S. Capitani, Nucl. Phys. **B592**, 18 (2001), hep-lat/0005008.
7. P. Hernández, K. Jansen, L. Lellouch, and H. Wittig, hep-lat/0106011.
8. S.J. Dong, T. Draper, I. Horváth, F.X. Lee, K.F. Liu, and J.B. Zhang, hep-lat/0108020.
9. C. Bernard and M. Golterman, Phys. Rev. D **46**, 853, (1992).
10. S.J. Dong, T. Draper, I. Horváth, F.X. Lee and J.B. Zhang, these proceedings.
11. ALPHA Collaboration, Nucl. Phys. **B571**, 237 (2000).
12. W. Bardeen, A. Duncan, E. Eichten, N. Isgur, and H. Thacker, hep-lat/0106008.
13. S.J. Dong, T. Draper, I. Horváth, F.X. Lee, K.F. Liu and J.B. Zhang, under preparation.